arXiv:0806.0967v1 [quant-ph] 5 Jun 2008

Gravitation of finite range and the accelerated expansion of the universe.

Bo E. Sernelius
Division of Theory and Modeling, Department of Physics,
Chemistry and Biology, Linköping University, SE-581 83 Linköping, Sweden[*]

In an earlier work [1] we showed that the gravitational interaction could be reproduced as a retarded dispersion interaction (Casimir interaction) between particles composed of hypothetical particles having harmonic oscillator interactions. Here we derive the modification of the force when the vacuum modes have a small but finite temperature. The resulting interaction is of finite range. We give the result on analytical form.

PACS numbers: 04.50.-h, 04.90.+e, 12.90.+b, 14.80.-j, 34.20.-b

In a previous work [1] we opened up for the possibility that gravity, considered to be one of the fundamental interactions in nature, is not a fundamental interaction after all. In our model, gravity is a dispersion interaction derived from more fundamental forces between particles having harmonic oscillator interactions. The dispersion forces are obtained from the change in the zero-point energy of the normal modes of the interaction fields in the system when the position of the composite particles are changed. This is in complete analogy with the electromagnetic dispersion forces between atoms [2, 3]. In the electromagnetic case there are two regions: the van der Waals region for moderate distances; the Casimir region for larger distances. We obtain corresponding regions in our model but with the proper choice of characteristic frequency of the composite particle the van der Waals region will never be observed; the range is smaller than the diameter of the composite particle. The dispersion force will have the r^{-2} dependence for all distances just as the gravitational force has.

The reason we started looking for an alternative origin of gravity is two-fold. One reason is that one has not been able to verify the existence of the graviton, the quantum of the gravitational field. The second is that recent experiments [4] have pointed to an accelerated expansion of the universe. "Vacuum energy is both the most plausible explanation and the most puzzling possibility", according to Ref. [4]. Another possible explanation could be found in a modified gravity. In the present work we pursue along this second path.

Big Bang occurred 13.7 billion years ago. Soon after the formation of atoms, matter and electromagnetic radiation decoupled. Since then the cosmic background radiation has expanded, in the same way as the rest of the universe, with a factor of approximately 11 000. All wave lengths have increased with, and the temperature of the cosmic background radiation has decreased with, this factor. The present temperature has been found to be 2.7 K. In analogy to this the radiation associated with the interaction between the fundamental particles in our model decoupled from matter at around the time when the composite particles were formed. This happened much earlier than the time of atom formation. Consequently the temperature of this cosmic background radiation is much lower than that of the electromagnetic counterpart.

Here we will derive the dispersion force at finite temperature and demonstrate that this force has a finite range. There is a temperature correction factor that goes towards zero when the distance goes towards infinity. The normal modes involved in the dispersion forces are of two types. One type is bound to the particles, the atoms in the electromagnetic case and the composite particles in the case of our model; it affects the forces only at close range, in the van der Waals range. The other type is a vacuum mode, modified by the boundary conditions set up by the presence of the atoms or composite particles; this type of mode dominates at large separations, in the Casimir range; it was the only type of mode in the classical work [5] by Casimir on the attraction between two perfectly conducting plates. This second type should be populated in accordance with the temperature of the cosmic background radiation. The first type, which we are not concerned with here, could be populated differently.

We found [1] the interaction potential at zero temperature is

$$\begin{aligned} V\left(r\right) &= -\tfrac{\hbar}{2\pi}\int\limits_0^\infty d\omega\alpha_1\left(i\omega\right)\alpha_2\left(i\omega\right)e^{-2\omega r/c}\left[3-6\left(\omega r/c\right)\right.\\ &\qquad \left.+\tfrac{17}{2}\left(\omega r/c\right)^2-\tfrac{9}{2}\left(\omega r/c\right)^3+\tfrac{3}{2}\left(\omega r/c\right)^4\right]\\ &= -\tfrac{\hbar}{2\pi}\alpha_1\left(0\right)\alpha_2\left(0\right)\int\limits_0^\infty d\omega e^{-2\omega r/c}\left[3-6\left(\omega r/c\right)\right.\\ &\qquad \left.+\tfrac{17}{2}\left(\omega r/c\right)^2-\tfrac{9}{2}\left(\omega r/c\right)^3+\tfrac{3}{2}\left(\omega r/c\right)^4\right], \end{aligned} \tag{1}$$

where we have used the fact that the exponential factor drops off so fast that only the static polarizabilities enter and can be brought outside the integral. Making the replacement $\alpha_i\left(0\right)=m_i\sqrt{32\pi\gamma/25\hbar c}; i=1,2$ of the static polarizabilities gives

$$\begin{aligned} V\left(r\right) &= -\gamma m_1 m_2 \left(\tfrac{4}{5}\right)^2 \tfrac{1}{c} \int\limits_0^\infty d\omega e^{-2\omega r/c} \left[3 - 6\left(\omega r/c\right) \right. \\ &\quad \left. + \tfrac{17}{2}\left(\omega r/c\right)^2 - \tfrac{9}{2}\left(\omega r/c\right)^3 + \tfrac{3}{2}\left(\omega r/c\right)^4\right] \\ &= -\gamma m_1 m_2 / r, \end{aligned} \tag{2}$$

where γ is the gravitational constant. The force is obtained as

$$\begin{aligned} F\left(r\right) &= -\tfrac{\partial V}{\partial r} \\ &= -\gamma m_1 m_2 \left(\tfrac{4}{5}\right)^2 \tfrac{1}{cr} \int\limits_0^\infty d\omega e^{-2\omega r/c} \left[-12\left(\omega r/c\right) \right. \\ &\quad + 29\left(\omega r/c\right)^2 - \tfrac{61}{2}\left(\omega r/c\right)^3 \\ &\quad \left. + 15\left(\omega r/c\right)^4 - 3\left(\omega r/c\right)^5\right] \\ &= -\gamma m_1 m_2 / r^2. \end{aligned} \tag{3}$$

At finite temperature the integral turns into a summation [3]

$$\int\limits_0^\infty d\omega \to \frac{2\pi}{\hbar\beta}\sum_{n=0}^{\infty}; \quad \omega_n = \frac{2\pi n}{\hbar\beta}, \tag{4}$$

where the $n = 0$ term should be multiplied by a factor of one half. In the present case we do not have to mind this since this term vanishes. We have

$$\begin{aligned} F\left(r\right) &= -\gamma m_1 m_2 \left(\tfrac{4}{5}\right)^2 \tfrac{2\pi}{\hbar\beta c r} \sum_{n=0}^{\infty} e^{-2\omega_n r/c} \left[-12\left(\omega_n r/c\right) \right. \\ &\quad + 29\left(\omega_n r/c\right)^2 - \tfrac{61}{2}\left(\omega_n r/c\right)^3 \\ &\quad \left. + 15\left(\omega_n r/c\right)^4 - 3\left(\omega_n r/c\right)^5\right]. \end{aligned} \tag{5}$$

This summation may be performed analytically and we find the temperature correction factor is

$$\begin{aligned} G\left(r,T\right) &= F\left(r,T\right)/F\left(r,0\right) \\ &= \left(\tfrac{4}{5}\right)^2 x \left[-12z^2 + 29z^3\left(x+1\right) \right. \\ &\quad - \tfrac{61}{2}z^4\left(x^2+4x+1\right) \\ &\quad + 15z^5\left(x^3+11x^2+11x+1\right) \\ &\quad \left. - 3z^6\left(x^4+26x^3+66x^2+26x+1\right)\right], \end{aligned} \tag{6}$$

where $y = 2\pi r/\hbar\beta c$, $x = \exp\left(-2y\right)$, $z = y/(1-x)$, and $\beta = 1/k_B T$. This temperature correction factor is displayed in Fig. (1).

In summary, we have demonstrated that the gravitational interaction derived from particles having harmonic oscillator interaction potentials has a finite range if the vacuum modes have a non-zero temperature. We have given the temperature correction factor on analytical form. This correction factor should have an effect on the expansion of the universe. To be noted is that the r and T dependences of the correction factor is in the form of the variable y. This means that the cut-off distance for gravity increases with the same speed as the expansion of the universe.



Figure 1: The temperature correction factor. See the text for details

This research was sponsored by EU within the EC-contract No:012142-NANOCASE and support from the VR Linné Centre LiLi-NFM and from CTS is gratefully acknowledged.

---

[*] Electronic address: bos@ifm.liu.se

[1] Bo E. Sernelius, *Gravitation as a Casimir interaction*, arXiv:0804.3054v1 [quant-ph] (2008).
[2] H. B. G. Casimir, and D. Polder, *Phys. Rev.* **73**, 360 (1948).
[3] Bo E. Sernelius, *Surface Modes in Physics*, Wiley-VCH (2001).
[4] M. S. Turner, and D. Huterer, *J. Phys. Soc. Jap.* **76**, 1110151 (2007).
[5] H. B. G. Casimir, *Proc. Koninklijke Nederlandse Akad. Wetenschappen* B **51**, 793-795 (1948).